

Logarithmic Functions - Important bases

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions. We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Our number system is base $b = 10$, so a commonly used log is: $\log_{10}(x)$

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Our number system is base $b = 10$, so a commonly used log is: $\log_{10}(x)$

In fact it is so commonly used that it is called the *common log*

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Our number system is base $b = 10$, so a commonly used log is: $\log_{10}(x)$

In fact it is so commonly used that it is called the *common log*

$$\log_{10}(x) = \log(x)$$

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Our number system is base $b = 10$, so a commonly used log is: $\log_{10}(x)$

In fact it is so commonly used that it is called the *common log*

$$\log_{10}(x) = \log(x)$$

Another often used base is base $b = e$ which came up in ► Our banking example

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Our number system is base $b = 10$, so a commonly used log is: $\log_{10}(x)$

In fact it is so commonly used that it is called the *common log*

$$\log_{10}(x) = \log(x)$$

Another often used base is base $b = e$ which came up in ► Our banking example

The logarithm with base $b = e$ is called the natural log and written as:

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Our number system is base $b = 10$, so a commonly used log is: $\log_{10}(x)$

In fact it is so commonly used that it is called the *common log*

$$\log_{10}(x) = \log(x)$$

Another often used base is base $b = e$ which came up in ► Our banking example

The logarithm with base $b = e$ is called the natural log and written as:

$$\log_e(x) = \ln(x)$$

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Our number system is base $b = 10$, so a commonly used log is: $\log_{10}(x)$

In fact it is so commonly used that it is called the *common log*

$$\log_{10}(x) = \log(x)$$

Another often used base is base $b = e$ which came up in ► Our banking example

The logarithm with base $b = e$ is called the natural log and written as:

$$\log_e(x) = \ln(x)$$

The natural log, $\ln(x)$, is the most frequently used logarithm

Logarithmic Functions - Important bases

► We defined Logarithms as the inverse functions of Exponential Functions.

We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Our number system is base $b = 10$, so a commonly used log is: $\log_{10}(x)$

In fact it is so commonly used that it is called the *common log*

$$\log_{10}(x) = \log(x)$$

Another often used base is base $b = e$ which came up in ► Our banking example

The logarithm with base $b = e$ is called the natural log and written as:

$$\log_e(x) = \ln(x)$$

The natural log, $\ln(x)$, is the most frequently used logarithm

We will wait to see why it is the "natural" choice in a Calculus course.

Logarithmic Functions - Important bases

► We defined

Logarithms as the inverse functions of Exponential Functions. We saw that, since there is a different exponential function for each base = b , there is a different logarithm for each base = b

For the function $y = f(x) = b^x$ the Logarithm base = b is:

$$f^{-1}(x) = \log_b(x)$$

Since $f^{-1}(x) = \log_b(x)$ is the inverse of $y = f(x) = b^x$ we have:

$$\log_b(b^x) = x \text{ and } b^{\log_b(x)} = x$$

Although, there is a different log for every base = b , some are more commonly used than others.

Our number system is base $b = 10$, so a commonly used log is: $\log_{10}(x)$

In fact it is so commonly used that it is called the *common log*

$$\log_{10}(x) = \log(x)$$

Another often used base is base $b = e$ which came up in ► Our banking example

The logarithm with base $b = e$ is called the natural log and written as:

$$\log_e(x) = \ln(x)$$

The natural log, $\ln(x)$, is the most frequently used logarithm

We will wait to see why it is the "natural" choice in a Calculus course.

Note: One other base that comes up frequently, typically in computer science, is base $b = 2$.