

# Reducing Fractions

# Reducing Fractions

► We found

that:

$$\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$$

and

$$\frac{n}{n} = 1 \text{ for } n \neq 0$$

# Reducing Fractions

► We found

that:

$$\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$$

and

$$\frac{n}{n} = 1 \text{ for } n \neq 0$$

Because of this, we can write one number in multiple different ways!

# Reducing Fractions

► We found that:  $\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$  and  $\frac{n}{n} = 1$  for  $n \neq 0$

Because of this, we can write one number in multiple different ways!

**Example:**

$$\frac{2}{6} =$$

# Reducing Fractions

► We found that:  $\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$  and  $\frac{n}{n} = 1$  for  $n \neq 0$

Because of this, we can write one number in multiple different ways!

**Example:**

$$\frac{2}{6} = \frac{2 \cdot 1}{2 \cdot 3}$$

# Reducing Fractions

► We found that:  $\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$  and  $\frac{n}{n} = 1$  for  $n \neq 0$

Because of this, we can write one number in multiple different ways!

**Example:**

$$\frac{2}{6} = \frac{2 \cdot 1}{2 \cdot 3} = \frac{2}{2} \cdot \frac{1}{3}$$

# Reducing Fractions

► We found that:  $\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$  and  $\frac{n}{n} = 1$  for  $n \neq 0$

Because of this, we can write one number in multiple different ways!

**Example:**

$$\frac{2}{6} = \frac{2 \cdot 1}{2 \cdot 3} = \underbrace{\frac{2}{2}}_1 \cdot \frac{1}{3}$$

# Reducing Fractions

► We found that:  $\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$  and  $\frac{n}{n} = 1$  for  $n \neq 0$

Because of this, we can write one number in multiple different ways!

**Example:**

$$\frac{2}{6} = \frac{2 \cdot 1}{2 \cdot 3} = \frac{\cancel{2}}{\underbrace{2}_{1}} \cdot \frac{1}{3} = \frac{1}{3}$$



# Reducing Fractions

► We found that:  $\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$  and  $\frac{n}{n} = 1$  for  $n \neq 0$

Because of this, we can write one number in multiple different ways!

**Example:**

$$\frac{2}{6} = \frac{2 \cdot 1}{2 \cdot 3} = \underbrace{\frac{\cancel{2}}{\cancel{2}}}_{1} \cdot \frac{1}{3} = \frac{1}{3}$$

So we have that:  $\frac{2}{6} = \frac{1}{3}$

# Reducing Fractions

► We found that:  $\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$  and  $\frac{n}{n} = 1$  for  $n \neq 0$

Because of this, we can write one number in multiple different ways!

**Example:**

$$\frac{2}{6} = \frac{2 \cdot 1}{2 \cdot 3} = \frac{\cancel{2}}{\underbrace{2}_{1}} \cdot \frac{1}{3} = \frac{1}{3}$$

So we have that:  $\frac{2}{6} = \frac{1}{3}$

This process re-writing a fraction with smaller numbers on top and bottom is called *reducing a fraction*

# Reducing Fractions

► We found that:  $\frac{a}{m} \cdot \frac{b}{n} = \frac{a \cdot b}{m \cdot n}$  and  $\frac{n}{n} = 1$  for  $n \neq 0$

Because of this, we can write one number in multiple different ways!

**Example:**

$$\frac{2}{6} = \frac{2 \cdot 1}{2 \cdot 3} = \underbrace{\frac{\cancel{2}}{\cancel{2}}}_{1} \cdot \frac{1}{3} = \frac{1}{3}$$

So we have that:  $\frac{2}{6} = \frac{1}{3}$

This process re-writing a fraction with smaller numbers on top and bottom is called *reducing a fraction*

Note: To reduce a fraction, there must be a common factor in the top and bottom that can be reduced to 1