

# Evaluating Basic Functions

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say " $y$  is a function of  $x$ "

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say "y is a function of x"

We write "y is a function of x" as:  $y = f(x)$

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say "y is a function of x"

We write "y is a function of x" as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it?

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say "y is a function of x"

We write "y is a function of x" as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it? In this class, we will look at mathematical formulas that tell how to find the  $y$ -value for each  $x$ -value

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say "y is a function of x"

We write "y is a function of x" as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it?

In this class, we will look at mathematical formulas that tell how to find the  $y$ -value for each  $x$ -value

In fact, we've already done this without calling them functions:



## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say "y is a function of x"

We write "y is a function of x" as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it?

In this class, we will look at mathematical formulas that tell how to find the  $y$ -value for each  $x$ -value

In fact, we've already done this without calling them functions:

$$y = -2x + 4$$

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say "y is a function of x"

We write "y is a function of x" as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it?

In this class, we will look at mathematical formulas that tell how to find the  $y$ -value for each  $x$ -value

In fact, we've already done this without calling them functions:

$$y = -2x + 4$$

Since each  $x$ -value gave one  $y$ -value, this was a function.

Although, we did not call them functions at the time.

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say " $y$  is a function of  $x$ "

We write " $y$  is a function of  $x$ " as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it?

In this class, we will look at mathematical formulas that tell how to find the  $y$ -value for each  $x$ -value

In fact, we've already done this without calling them functions:

$$y = -2x + 4$$

Since each  $x$ -value gave one  $y$ -value, this was a function.

Although, we did not call them functions at the time.

We were able to find those  $y$ -values by picking an  $x$ -value and evaluating the right hand side to find the  $y$ -value.

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say " $y$  is a function of  $x$ "

We write " $y$  is a function of  $x$ " as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it?

In this class, we will look at mathematical formulas that tell how to find the  $y$ -value for each  $x$ -value

In fact, we've already done this without calling them functions:

$$y = -2x + 4$$

Since each  $x$ -value gave one  $y$ -value, this was a function.

Although, we did not call them functions at the time.

We were able to find those  $y$ -values by picking an  $x$ -value and evaluating the right hand side to find the  $y$ -value.

Before: if  $x = 1$  then:  $y = -2 \cdot 1 + 4 = -2 + 4 = 2$ ; sol:  $(1, 2)$

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say " $y$  is a function of  $x$ "

We write " $y$  is a function of  $x$ " as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it?

In this class, we will look at mathematical formulas that tell how to find the  $y$ -value for each  $x$ -value

In fact, we've already done this without calling them functions:

$$y = -2x + 4$$

Since each  $x$ -value gave one  $y$ -value, this was a function.

Although, we did not call them functions at the time.

We were able to find those  $y$ -values by picking an  $x$ -value and evaluating the right hand side to find the  $y$ -value.

Before: if  $x = 1$  then:  $y = -2 \cdot 1 + 4 = -2 + 4 = 2$ ; sol:  $(1, 2)$

Now:  $y = f(1) = -2 \cdot 1 + 4 = -2 + 4 = 2$ ;  $f(1) = 2$

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say " $y$  is a function of  $x$ "

We write " $y$  is a function of  $x$ " as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it?

In this class, we will look at mathematical formulas that tell how to find the  $y$ -value for each  $x$ -value

In fact, we've already done this without calling them functions:

$$y = -2x + 4$$

Since each  $x$ -value gave one  $y$ -value, this was a function.

Although, we did not call them functions at the time.

We were able to find those  $y$ -values by picking an  $x$ -value and evaluating the right hand side to find the  $y$ -value.

Before: if  $x = 1$  then:  $y = -2 \cdot 1 + 4 = -2 + 4 = 2$ ; sol:  $(1, 2)$

Now:  $y = f(1) = -2 \cdot 1 + 4 = -2 + 4 = 2$ ;  $f(1) = 2$

This is called evaluating the function.

## Evaluating Basic Functions

We saw that functions are sets of ordered pairs of numbers  $(x, y)$  so that no two ordered pairs have the same  $x$ -value. This means that if we know the  $x$ -value of the ordered pair, there can be just one  $y$ -value.

So, the  $y$ -value of the pair depends on the  $x$ -value, which leads us to say " $y$  is a function of  $x$ "

We write " $y$  is a function of  $x$ " as:  $y = f(x)$

There is just one  $y$ -value for each  $x$ -value but what is it?

In this class, we will look at mathematical formulas that tell how to find the  $y$ -value for each  $x$ -value

In fact, we've already done this without calling them functions:

$$y = -2x + 4$$

Since each  $x$ -value gave one  $y$ -value, this was a function.

Although, we did not call them functions at the time.

We were able to find those  $y$ -values by picking an  $x$ -value and evaluating the right hand side to find the  $y$ -value.

Before: if  $x = 1$  then:  $y = -2 \cdot 1 + 4 = -2 + 4 = 2$ ; sol:  $(1, 2)$

Now:  $y = f(1) = -2 \cdot 1 + 4 = -2 + 4 = 2$ ;  $f(1) = 2$

This is called evaluating the function.

Functions make what we've already been doing more concise!