

# 1-1 Functions

# 1-1 Functions

► Recall: we define a *function* as a set of ordered pairs  $(x, y)$  so that no two pairs have the same  $x$ -value.

# 1-1 Functions

► Recall: we define a *function* as a set of ordered pairs  $(x, y)$  so that no two pairs have the same  $x$ -value.

In other words, for  $x$ -value there is only one  $y$ -value.

# 1-1 Functions

► Recall: we define a *function* as a set of ordered pairs  $(x, y)$  so that no two pairs have the same  $x$ -value.

In other words, for  $x$ -value there is only one  $y$ -value.

However, a function can have one  $y$ - paired with two  $x$ -values.

## 1-1 Functions

► Recall: we define a *function* as a set of ordered pairs  $(x, y)$  so that no two pairs have the same  $x$ -value.

In other words, for  $x$ -value there is only one  $y$ -value.

However, a function can have one  $y$ - paired with two  $x$ -values.

For example,  $f(x) = x^2$  is a function even though  $(3, 9)$  and  $(-3, 9)$  are two points with the same  $y$ -value.

# 1-1 Functions

► Recall: we define a *function* as a set of ordered pairs  $(x, y)$  so that no two pairs have the same  $x$ -value.

In other words, for  $x$ -value there is only one  $y$ -value.

However, a function can have one  $y$ - paired with two  $x$ -values.

For example,  $f(x) = x^2$  is a function even though  $(3, 9)$  and  $(-3, 9)$  are two points with the same  $y$ -value.

For some functions though, there are no two pairs that have the same  $y$ -value.

## 1-1 Functions

► Recall: we define a *function* as a set of ordered pairs  $(x, y)$  so that no two pairs have the same  $x$ -value.

In other words, for  $x$ -value there is only one  $y$ -value.

However, a function can have one  $y$ - paired with two  $x$ -values.

For example,  $f(x) = x^2$  is a function even though  $(3, 9)$  and  $(-3, 9)$  are two points with the same  $y$ -value.

For some functions though, there are no two pairs that have the same  $y$ -value.

Which means that for these special functions there is exactly 1  $x$ -value for each  $y$ -value, and exactly 1  $y$ -value for each  $x$ -value.

# 1-1 Functions

► Recall: we define a *function* as a set of ordered pairs  $(x, y)$  so that no two pairs have the same  $x$ -value.

In other words, for  $x$ -value there is only one  $y$ -value.

However, a function can have one  $y$ - paired with two  $x$ -values.

For example,  $f(x) = x^2$  is a function even though  $(3, 9)$  and  $(-3, 9)$  are two points with the same  $y$ -value.

For some functions though, there are no two pairs that have the same  $y$ -value.

Which means that for these special functions there is exactly 1  $x$ -value for each  $y$ -value, and exactly 1  $y$ -value for each  $x$ -value.

These functions are called 1-1 (*one-to-one*) functions