

Intro to Graphing Polynomials

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_2 x^2 + a_1 x + a_0$$

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y -  by finding the point when $x = 0$

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y -  by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \dots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0$$

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y -  by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \dots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y -  by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \dots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

So, for any polynomial, the y -intercept is: $(0, a_0)$

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y -  by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \dots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

So, for any polynomial, the y -intercept is: $(0, a_0)$

What about the x -intercepts?

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y -  by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \dots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

So, for any polynomial, the y -intercept is: $(0, a_0)$

What about the x -intercepts? $y = 0$

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials.

Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y -  by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \dots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

So, for any polynomial, the y -intercept is: $(0, a_0)$

What about the x -intercepts? $y = 0$

So, to find the x -intercepts we need to solve:

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials. Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y -  by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \dots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

So, for any polynomial, the y -intercept is: $(0, a_0)$

What about the x -intercepts? $y = 0$

So, to find the x -intercepts we need to solve:

$$0 = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials. Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y -  by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \dots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

So, for any polynomial, the y -intercept is: $(0, a_0)$

What about the x -intercepts? $y = 0$

So, to find the x -intercepts we need to solve:

$$0 = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

There is no perfect way to solve this.

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials. Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

We can find the y - [intercept](#) by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \dots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

So, for any polynomial, the y -intercept is: $(0, a_0)$

What about the x -intercepts? $y = 0$

So, to find the x -intercepts we need to solve:

$$0 = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

There is no perfect way to solve this.

Our best method is to find the roots is with the [Rational Root Theorem](#)

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials. Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_2 x^2 + a_1 x + a_0$$

We can find the y - [intercept](#) by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \cdots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

So, for any polynomial, the y -intercept is: $(0, a_0)$

What about the x -intercepts? $y = 0$

So, to find the x -intercepts we need to solve:

$$0 = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_2 x^2 + a_1 x + a_0$$

There is no perfect way to solve this.

Our best method is to find the roots is with the [Rational Root Theorem](#)

Once we find the roots $x = k_1, k_2, k_3, \dots$ then the x -intercepts are:

Intro to Graphing Polynomials

When graphing lines and quadratics, we used the x - and y -intercepts as important reference points in our graphs.

Similarly, the x - and y -int are important in graphing Polynomials. Also similar to linear and quadratic functions, it will be easier to find the y -intercept than the x -intercept.

For a polynomial:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_2 x^2 + a_1 x + a_0$$

We can find the y - [intercept](#) by finding the point when $x = 0$

$$P(0) = a_n \cdot 0^n + a_{n-1} \cdot 0^{n-1} + \cdots + a_2 \cdot 0^2 + a_1 \cdot 0 + a_0 = a_0$$

So, for any polynomial, the y -intercept is: $(0, a_0)$

What about the x -intercepts? $y = 0$

So, to find the x -intercepts we need to solve:

$$0 = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_2 x^2 + a_1 x + a_0$$

There is no perfect way to solve this.

Our best method is to find the roots is with the [Rational Root Theorem](#)

Once we find the roots $x = k_1, k_2, k_3, \dots$ then the x -intercepts are:


$$(k_1, 0), (k_2, 0), (k_3, 0), \dots$$

Intro to Graphing Polynomials

Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:


x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$



Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$




Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?




Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.



Intro to Graphing Polynomials


Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.

The graph does not change from pos. to neg. (or vice-versa).



Intro to Graphing Polynomials


Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.

The graph does not change from pos. to neg. (or vice-versa). Why not?



Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:


x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.

The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them



Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:


x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.

The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them



Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$


What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.

The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them

This is the same between each pair of x -intercepts



Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?


We might not know exactly what the graph looks like.

The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them

This is the same between each pair of x -intercepts

Notice: For our graph to connect the x -int there are *Turning Points*



Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.


The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them

This is the same between each pair of x -intercepts

Notice: For our graph to connect the x -int there are *Turning Points*

We will need to wait until Calculus to pinpoint the coords of *Turning Points*



Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.


The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them

This is the same between each pair of x -intercepts

Notice: For our graph to connect the x -int there are *Turning Points*

We will need to wait until Calculus to pinpoint the coords of *Turning Points*



What remains to understand is what happens at the ends of the graph.

Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.


The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them

This is the same between each pair of x -intercepts

Notice: For our graph to connect the x -int there are *Turning Points*

We will need to wait until Calculus to pinpoint the coords of *Turning Points*



What remains to understand is what happens at the ends of the graph.

Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.


The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them

This is the same between each pair of x -intercepts

Notice: For our graph to connect the x -int there are *Turning Points*

We will need to wait until Calculus to pinpoint the coords of *Turning Points*



What remains to understand is what happens at the ends of the graph.

Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.


The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them

This is the same between each pair of x -intercepts

Notice: For our graph to connect the x -int there are *Turning Points*

We will need to wait until Calculus to pinpoint the coords of *Turning Points*



What remains to understand is what happens at the ends of the graph.

Intro to Graphing Polynomials

Suppose that we find **all** of the x - and y -intercepts are:

x -int: $(k_1, 0), (k_2, 0), (k_3, 0)$ and y -int: $(0, a_0)$

What does the graph look like between the intercepts?

We might not know exactly what the graph looks like.


The graph does not change from pos. to neg. (or vice-versa). Why not?

Because then there would need to be another x -intercept between them

This is the same between each pair of x -intercepts

Notice: For our graph to connect the x -int there are *Turning Points*

We will need to wait until Calculus to pinpoint the coords of *Turning Points*



What remains to understand is what happens at the ends of the graph. In other words, what happens for really large x -values?