So far, the two functions that we explored most deeply are:

So far, the two functions that we explored most deeply are: Linear functions: y = f(x) = mx + b

So far, the two functions that we explored most deeply are: Linear functions: y = f(x) = mx + bQuadratic functions: $y = f(x) = ax^2 + bx + c$

So far, the two functions that we explored most deeply are: Linear functions: Quadratic functions: y = f(x) = mx + b $y = f(x) = ax^2 + bx + c$ Notice the similarity between these.

So far, the two functions that we explored most deeply are: Linear functions: Quadratic functions:

y = f(x) = mx + bNotice the similarity between these. $y = f(x) = ax^2 + bx + c$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^1 + bx^0$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^2 + bx^1 + cx^0$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^1 + bx^0$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^2 + bx^1 + cx^0$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^1 + bx^0$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^2 + bx^1 + cx^0$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2

So, the next logical step is to see what happens if we include x^3

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^1 + bx^0$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^2 + bx^1 + cx^0$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^1 + bx^0$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^2 + bx^1 + cx^0$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4 And after that x^5 , x^6 , etc.

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^{1} + bx^{0}$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^{2} + bx^{1} + cx^{0}$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

And after that
$$x^5$$
, x^6 , etc.

We would need A LOT of time to talk about each one individually

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^{1} + bx^{0}$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^{2} + bx^{1} + cx^{0}$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

And after that
$$x^5$$
, x^6 , etc.

We would need A LOT of time to talk about each one individually So, let's study them all at once!

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^{1} + bx^{0}$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^{2} + bx^{1} + cx^{0}$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

And after that
$$x^5$$
, x^6 , etc.

We would need A LOT of time to talk about each one individually So, let's study them all at once!

A *Polynomial* is a function of the form:

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^{1} + bx^{0}$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^{2} + bx^{1} + cx^{0}$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

And after that x^5 , x^6 , etc.

We would need A LOT of time to talk about each one individually So, let's study them all at once!

A Polynomial is a function of the form:

$$y = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$$

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^{1} + bx^{0}$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^{2} + bx^{1} + cx^{0}$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

And after that x^5 , x^6 , etc.

We would need A LOT of time to talk about each one individually So, let's study them all at once!

A *Polynomial* is a function of the form:

 $y = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$ The highest power of x, in this case n, is called the *degree*

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^{1} + bx^{0}$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^{2} + bx^{1} + cx^{0}$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

And after that x^5 , x^6 , etc.

We would need A LOT of time to talk about each one individually So, let's study them all at once!

A *Polynomial* is a function of the form:

 $y = \mathbf{a_n}x^n + \mathbf{a_{n-1}}x^{n-1} + \dots + \mathbf{a_2}x^2 + \mathbf{a_1}x + \mathbf{a_0}$ The highest power of x, in this case n, is called the *degree* Because n can be big, we don't use a, b, c, etc. for the *coefficients*

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^{1} + bx^{0}$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^{2} + bx^{1} + cx^{0}$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

And after that x^5 , x^6 , etc.

We would need A LOT of time to talk about each one individually So, let's study them all at once!

A *Polynomial* is a function of the form:

 $y = \mathbf{a_n}x^n + \mathbf{a_{n-1}}x^{n-1} + \dots + \mathbf{a_2}x^2 + \mathbf{a_1}x + \mathbf{a_0}$ The highest power of x, in this case n, is called the *degree* Because n can be big, we don't use a, b, c, etc. for the *coefficients* The coefficient of x^n is $\mathbf{a_n}$ is called the *lead coefficient*.

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^1 + bx^0$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^2 + bx^1 + cx^0$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

And after that x^5 , x^6 , etc.

We would need A LOT of time to talk about each one individually So, let's study them all at once!

A *Polynomial* is a function of the form:

 $y = \mathbf{a_n}x^n + \mathbf{a_{n-1}}x^{n-1} + \dots + \mathbf{a_2}x^2 + \mathbf{a_1}x + \mathbf{a_0}$ The highest power of x, in this case n, is called the *degree* Because n can be big, we don't use a, b, c, etc. for the *coefficients* The coefficient of x^n is $\mathbf{a_n}$ is called the *lead coefficient*. And $\mathbf{a_n}x^n$ is called the *lead term*.

So far, the two functions that we explored most deeply are:

Linear functions: $y = f(x) = mx^{1} + bx^{0}$ Notice the similarity between these. Quadratic functions: $y = f(x) = ax^{2} + bx^{1} + cx^{0}$

Although it might not be obvious at first glance, in both of them we are adding up powers of x

The jump from linear to quadratic meant we included x^2 So, the next logical step is to see what happens if we include x^3 But then we'll need to include x^4

And after that x^5 , x^6 , etc.

We would need A LOT of time to talk about each one individually So, let's study them all at once!

A *Polynomial* is a function of the form:

 $P(x) = y = \mathbf{a_n}x^n + \mathbf{a_{n-1}}x^{n-1} + \dots + \mathbf{a_2}x^2 + \mathbf{a_1}x + \mathbf{a_0}$ The highest power of x, in this case n, is called the *degree* Because n can be big, we don't use a, b, c, etc. for the *coefficients* The coefficient of xⁿ is a_n is called the *lead coefficient*. And a_nxⁿ is called the *lead term*.