Recall: The Quadratic Formula says r_1 and r_2 are the roots of: $ax^2 + bx + c = 0$

Recall: The Quadratic Formula says r_1 and r_2 are the roots of: $ax^2 + bx + c = 0$ $r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$

Recall: The \bigcirc Quadratic Formula says r_1 and r_2 are \bigcirc the roots of:

$$r_{1,2} = \frac{ax^2 + bx + c = 0}{-(b) \pm \sqrt{(b)^2 - 4ac}}$$

Recall: The \bigcirc Quadratic Formula says r_1 and r_2 are \bigcirc the roots of:

$$r_{1,2} = \frac{ax^2 + bx + c = 0}{-(b) \pm \sqrt{(b)^2 - 4ac}}$$

$$r_{1,2} = \frac{-(4) \pm \sqrt{(4)^2 - 4 \cdot 1 \cdot 5}}{2 \cdot 1}$$

Recall: The Quadratic Formula says r_1 and r_2 are the roots of:

e Quadratic Formula says
$$r_1$$
 and r_2 are the roots of
$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(4) \pm \sqrt{(4)^2 - 4 \cdot 1 \cdot 5}}{2 \cdot 1}$$

$$-4 \pm \sqrt{16 - 20}$$

$$\frac{\sqrt{(4)^2 - 4 \cdot 1}}{2 \cdot 1}$$

$$\frac{2 \cdot 1}{16 - 20}$$

Recall: The Quadratic Formula says r_1 and r_2 are the roots of:

the Quadratic Formula says
$$r_1$$
 and r_2 are the roots
$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(4) \pm \sqrt{(4)^2 - 4 \cdot 1 \cdot 5}}{2 \cdot 1}$$

$$A + \sqrt{16 - 20}$$

$$r_{1,2} = \frac{-(4) \pm \sqrt{(4)^2 - 4 \cdot 1 \cdot 5}}{2 \cdot 1}$$
$$= \frac{-4 \pm \sqrt{16 - 20}}{2}$$

 $=\frac{-4\pm\sqrt{-4}}{2}$

Recall: The \bigcirc Quadratic Formula says r_1 and r_2 are \bigcirc the roots of:

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(4) \pm \sqrt{(4)^2 - 4 \cdot 1 \cdot 5}}{2 \cdot 1}$$

$$= \frac{-4 \pm \sqrt{16 - 20}}{2}$$

$$= \frac{-4 \pm \sqrt{-4}}{2}$$

Since -4 < 0 we have no real solutions.