Example: Find the Domain of:

$$f(x) = \sqrt{6x^2 + x - 2}$$

Example: Find the Domain of:

$$f(x) = \sqrt{6x^2 + x - 2}$$

Precall: The Domain of a function is the set of x values of the function.

Example: Find the Domain of:

$$f(x) = \sqrt{6x^2 + x - 2}$$

Precall: The Domain of a function is the set of x values of the function.

• We saw that the domain of f(x) consists of all x so that:

$$6x^2 + x - 2 \ge 0$$

Example: Find the Domain of:

$$f(x) = \sqrt{6x^2 + x - 2}$$

Precall: The Domain of a function is the set of x values of the function. **Precall:** We saw that the domain of f(x) consists of all x so that:

$$6x^2 + x - 2 \ge 0$$

Another way we can think about this is that f(x) is undefined if:

 $6x^2 + x - 2 \le 0$

Example: Find the Domain of: $f(x) = \sqrt{6x^2 + x - 2}$ Recall: The Domain of a function is the set of x values of the function. We saw that the domain of f(x) consists of all x so that: $6x^2 + x - 2 \ge 0$ Another way we can think about this is that f(x) is undefined if: $6x^2 + x - 2 \le 0$ We solve this inequality to find:

Example: Find the Domain of: $f(x) = \sqrt{6x^2 + x - 2}$ Recall: The Domain of a function is the set of x values of the function. We saw that the domain of f(x) consists of all x so that: $6x^2 + x - 2 \ge 0$ Another way we can think about this is that f(x) is undefined if: $6x^2 + x - 2 \le 0$ We solve this inequality to find: $\frac{-2}{-3} < x < \frac{1}{2}$

Example: Find the Domain of: $f(x) = \sqrt{6x^2 + x - 2}$ **P**Recall: The Domain of a function is the set of x values of the function. • We saw that the domain of f(x) consists of all x so that: $6x^2 + x - 2 > 0$ Another way we can think about this is that f(x) is undefined if: $6x^2 + x - 2 \le 0$ • We solve this inequality to find: $\frac{-2}{3} < x < \frac{1}{2}$ Since f(x) is not define for $\frac{-2}{3} < x < \frac{1}{2}$

The Domain is all other x-values.

Example: Find the Domain of: $f(x) = \sqrt{6x^2 + x - 2}$ **Precall:** The Domain of a function is the set of x values of the function. • We saw that the domain of f(x) consists of all x so that: $6x^2 + x - 2 > 0$ Another way we can think about this is that f(x) is undefined if: $6x^2 + x - 2 < 0$ • We solve this inequality to find: $\frac{-2}{3} < x < \frac{1}{2}$ Since f(x) is not define for $\frac{-2}{3} < x < \frac{1}{2}$ The Domain is all other x-values. **Conclusion:** Domain of $f(x) = \sqrt{6x^2 + x - 2}$ is: $\left(-\infty, \frac{-2}{3}\right] \cup \left|\frac{1}{2}, \infty\right)$

Example: Find the Domain of: $f(x) = \sqrt{6x^2 + x - 2}$ **P**Recall: The Domain of a function is the set of x values of the function. • We saw that the domain of f(x) consists of all x so that: $6x^2 + x - 2 > 0$ Another way we can think about this is that f(x) is undefined if: $6x^2 + x - 2 < 0$ • We solve this inequality to find: $\frac{-2}{3} < x < \frac{1}{2}$ Since f(x) is not define for $\frac{-2}{3} < x < \frac{1}{2}$ The Domain is all other x-values. **Conclusion:** Domain of $f(x) = \sqrt{6x^2 + x - 2}$ is: $\left(-\infty, \frac{-2}{3}\right] \cup \left|\frac{1}{2}, \infty\right)$ We can visualize the Domain on the number line:

Example: Find the Domain of: $f(x) = \sqrt{6x^2 + x - 2}$ **P**Recall: The Domain of a function is the set of x values of the function. • We saw that the domain of f(x) consists of all x so that: $6x^2 + x - 2 > 0$ Another way we can think about this is that f(x) is undefined if: $6x^2 + x - 2 \le 0$ • We solve this inequality to find: $\frac{-2}{3} < x < \frac{1}{2}$ Since f(x) is not define for $\frac{-2}{2} < x < \frac{1}{2}$ The Domain is all other x-values. **Conclusion:** Domain of $f(x) = \sqrt{6x^2 + x - 2}$ is: $\left(-\infty, \frac{-2}{3}\right] \cup \left|\frac{1}{2}, \infty\right)$ We can visualize the Domain on the number line:

