

Reducing Rational Functions

Reducing Rational Functions

► We define a *Rational Function* to be:

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} =$$

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} =$$

Like fractions, we will look for common factors in the top and bottom.

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} =$$

Like fractions, we will look for common factors in the top and bottom. To do this, we should factor the top to see what the factors are, and if there are any common factors with the bottom.

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} = \frac{3 \cdot (x + 2)}{3x}$$

Like fractions, we will look for common factors in the top and bottom. To do this, we should factor the top to see what the factors are, and if there are any common factors with the bottom.

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} = \frac{3 \cdot (x + 2)}{3x} = \frac{3}{3} \cdot \frac{x + 2}{x}$$

Like fractions, we will look for common factors in the top and bottom. To do this, we should factor the top to see what the factors are, and if there are any common factors with the bottom.

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} = \frac{3 \cdot (x + 2)}{3x} = \frac{\cancel{3}}{\cancel{3}} \cdot \frac{x + 2}{x} = \frac{x + 2}{x}$$

Like fractions, we will look for common factors in the top and bottom. To do this, we should factor the top to see what the factors are, and if there are any common factors with the bottom.

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} = \frac{3 \cdot (x + 2)}{3x} = \frac{\cancel{3}}{\cancel{3}} \cdot \frac{x + 2}{x} = \frac{x + 2}{x}$$

Like fractions, we will look for common factors in the top and bottom.

To do this, we should factor the top to see what the factors are, and if there are any common factors with the bottom.

Can we reduce this any more?

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} = \frac{3 \cdot (x + 2)}{3x} = \frac{\cancel{3}}{\cancel{3}} \cdot \frac{x + 2}{x} = \frac{x + 2}{x}$$

Like fractions, we will look for common factors in the top and bottom.

To do this, we should factor the top to see what the factors are, and if there are any common factors with the bottom.

Can we reduce this any more?

While we have a common term of x in the top and bottom, we can only reduce common factors in a fraction!

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} = \frac{3 \cdot (x + 2)}{3x} = \frac{\cancel{3}}{\cancel{3}} \cdot \frac{x + 2}{x} = \frac{x + 2}{x}$$

Like fractions, we will look for common factors in the top and bottom.

To do this, we should factor the top to see what the factors are, and if there are any common factors with the bottom.

Can we reduce this any more?

While we have a common term of x in the top and bottom, we can only reduce common factors in a fraction!

Our factor up top is: $(x + 2)$ while the bottom has the factor: x

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} = \frac{3 \cdot (x + 2)}{3x} = \frac{\cancel{3}}{\cancel{3}} \cdot \frac{x + 2}{x} = \frac{x + 2}{x}$$

Like fractions, we will look for common factors in the top and bottom.

To do this, we should factor the top to see what the factors are, and if there are any common factors with the bottom.

Can we reduce this any more?

While we have a common term of x in the top and bottom, we can only reduce common factors in a fraction!

Our factor up top is: $(x + 2)$ while the bottom has the factor: x

Since these are not the same, we cannot reduce further.

Reducing Rational Functions

► We define a *Rational Function* to be:

$$f(x) = \frac{P(x)}{D(x)}$$

Where $P(x)$ and $D(x)$ are both Polynomials.

► Like fractions: Rational expressions can be simplified.

We will use similar techniques to reduce Rationals as fractions.

Example: Simplify:

$$\frac{3x + 6}{3x} = \frac{3 \cdot (x + 2)}{3x} = \frac{\cancel{3}}{\cancel{3}} \cdot \frac{x + 2}{x} = \frac{x + 2}{x}$$

Like fractions, we will look for common factors in the top and bottom.

To do this, we should factor the top to see what the factors are, and if there are any common factors with the bottom.

Can we reduce this any more?

While we have a common term of x in the top and bottom, we can only reduce common factors in a fraction!

Our factor up top is: $(x + 2)$ while the bottom has the factor: x

Since these are not the same, we cannot reduce further.

It is as important to know when we cannot simplify as when we can!