

Changing Bases of Logarithms - Example 2

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Since we cannot compute \log_5 on our calculators, but can compute $\ln = \ln_e$, we want to change the base using our [Change of Base Formula](#)

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Since we cannot compute \log_5 on our calculators, but can compute $\ln = \ln_e$, we want to change the base using our [Change of Base Formula](#)

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}$$

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Since we cannot compute \log_5 on our calculators, but can compute $\ln = \ln_e$, we want to change the base using our [Change of Base Formula](#)

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}$$

Changing the base from 5 to e , we get:

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Since we cannot compute \log_5 on our calculators, but can compute $\ln = \ln_e$, we want to change the base using our [Change of Base Formula](#)

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}$$

Changing the base from 5 to e , we get:

$$\log_5(30) = \frac{\ln(30)}{\ln(5)}$$

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Since we cannot compute \log_5 on our calculators, but can compute $\ln = \ln_e$, we want to change the base using our [Change of Base Formula](#)

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}$$

Changing the base from 5 to e , we get:

$$\log_5(30) = \frac{\ln(30)}{\ln(5)}$$

We can compute the right side with a calculator using the \log button

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Since we cannot compute \log_5 on our calculators, but can compute $\ln = \ln_e$, we want to change the base using our [Change of Base Formula](#)

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}$$

Changing the base from 5 to e , we get:

$$\log_5(30) = \frac{\ln(30)}{\ln(5)} \approx \frac{3.4011973}{1.6094379}$$

We can compute the right side with a calculator using the \log button

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Since we cannot compute \log_5 on our calculators, but can compute $\ln = \ln_e$, we want to change the base using our [Change of Base Formula](#)

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}$$

Changing the base from 5 to e , we get:

$$\log_5(30) = \frac{\ln(30)}{\ln(5)} \approx \frac{3.4011973}{1.6094379} \approx 2.113$$

We can compute the right side with a calculator using the \log button

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Since we cannot compute \log_5 on our calculators, but can compute $\ln = \ln_e$, we want to change the base using our [Change of Base Formula](#)

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}$$

Changing the base from 5 to e , we get:

$$\log_5(30) = \frac{\ln(30)}{\ln(5)} \approx \frac{3.4011973}{1.6094379} \approx 2.113$$

We can compute the right side with a calculator using the \log button

Conclusion: $\log_5(30) \approx 2.113$

Changing Bases of Logarithms - Example 2

Example: Compute $\log_5(30)$ rounded to 3 decimal places

Since we cannot compute \log_5 on our calculators, but can compute $\ln = \ln_e$, we want to change the base using our [Change of Base Formula](#)

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}$$

Changing the base from 5 to e , we get:

$$\log_5(30) = \frac{\ln(30)}{\ln(5)} \approx \frac{3.4011973}{1.6094379} \approx 2.113$$

We can compute the right side with a calculator using the \log button

Conclusion: $\log_5(30) \approx 2.113$

Note: We could have changed the base to either 10 or e but had to pick one.