

Solving Quadratic Equations with Factoring - Example 2

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Factors of 3 are:

$$1 \cdot 3 = 3$$

$$-1 \cdot (-3) = 3$$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Factors of 3 are:

$$1 \cdot 3 = 3 \text{ and } 1 + 3 = 4$$

$$-1 \cdot (-3) = 3$$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Factors of 3 are:

$$1 \cdot 3 = 3 \text{ and } 1 + 3 = 4$$

$$-1 \cdot (-3) = 3$$

$$s, t = 1, 3$$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Factors of 3 are:

$$1 \cdot 3 = 3 \text{ and } 1 + 3 = 4$$

$$-1 \cdot (-3) = 3$$

$$s, t = 1, 3$$

So, we can write $x^2 + 4x + 3$ as:

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Factors of 3 are:

$$1 \cdot 3 = 3 \text{ and } 1 + 3 = 4$$

$$-1 \cdot (-3) = 3$$

$$s, t = 1, 3$$

So, we can write $x^2 + 4x + 3$ as:

$$x^2 + 4x + 3 = (x + 1) \cdot (x + 3)$$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Factors of 3 are:

$$1 \cdot 3 = 3 \text{ and } 1 + 3 = 4$$

$$-1 \cdot (-3) = 3$$

$$s, t = 1, 3$$

So, we can write $x^2 + 4x + 3$ as:

$$x^2 + 4x + 3 = (x + 1) \cdot (x + 3)$$

Remember, we want to solve:

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Factors of 3 are:

$$1 \cdot 3 = 3 \text{ and } 1 + 3 = 4$$

$$-1 \cdot (-3) = 3$$

$$s, t = 1, 3$$

So, we can write $x^2 + 4x + 3$ as:

$$x^2 + 4x + 3 = (x + 1) \cdot (x + 3)$$

Remember, we want to solve:

$$0 = x^2 + 4x + 3 = (x + 1) \cdot (x + 3)$$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Factors of 3 are:

$$1 \cdot 3 = 3 \text{ and } 1 + 3 = 4$$

$$-1 \cdot (-3) = 3$$

$$s, t = 1, 3$$

So, we can write $x^2 + 4x + 3$ as:

$$x^2 + 4x + 3 = (x + 1) \cdot (x + 3)$$

Remember, we want to solve:

$$0 = x^2 + 4x + 3 = (x + 1) \cdot (x + 3)$$

Using ▶ The Zero Product Property we get $[x + 1 = 0 \text{ or } x + 3 = 0$

Solving Quadratic Equations with Factoring - Example 2

Example: Find solutions to:

$$x^2 + 4x + 3 = 0$$

To solve this, we want to write $x^2 + 4x + 3 = (x + s) \cdot (x + t)$

This is called factored form.

We need to find what s and t are.

▶ As before we distribute the right hand side to get:

$$x^2 + 4x + 3 = (x + s) \cdot (x + t) = x^2 + (s + t)x + st$$

To make sure both sides are the same, we need $s + t = 4$ and $s \cdot t = 3$

Let's start by thinking of all pairs of integers s and t can be multiplied together so: $s \cdot t = 3$

Factors of 3 are:

$$1 \cdot 3 = 3 \text{ and } 1 + 3 = 4$$

$$-1 \cdot (-3) = 3$$

$$s, t = 1, 3$$

So, we can write $x^2 + 4x + 3$ as:

$$x^2 + 4x + 3 = (x + 1) \cdot (x + 3)$$

Remember, we want to solve:

$$0 = x^2 + 4x + 3 = (x + 1) \cdot (x + 3)$$

Using ▶ The Zero Product Property we get $[x + 1 = 0 \text{ or } x + 3 = 0$

The solutions to $0 = x^2 + 4x + 3$ are $x = -1, -3$