We will start our discussion of Quadratic Equations with an example

We will start our discussion of Quadratic Equations with an example **Example:** Find the solution(s) to:

 $(x-3)\cdot(x+2)=0$

We will start our discussion of Quadratic Equations with an example **Example:** Find the solution(s) to:

$$(x-3)\cdot(x+2)=0$$

Here, we have two numbers multiplied together to get $\ensuremath{0}$

We will start our discussion of Quadratic Equations with an example **Example:** Find the solution(s) to:

 $(x-3)\cdot(x+2)=0$

Here, we have two numbers multiplied together to get 0 The two numbers we multiply together are (x - 3) and (x + 2)

We will start our discussion of Quadratic Equations with an example **Example:** Find the solution(s) to:

 $(x-3)\cdot(x+2)=0$

Here, we have two numbers multiplied together to get 0 The two numbers we multiply together are (x - 3) and (x + 2)

• By the Zero-Product Property either:

$$(x-3)=0$$

$$(x+2)=0$$

We will start our discussion of Quadratic Equations with an example **Example:** Find the solution(s) to:

 $(x-3)\cdot(x+2)=0$

Here, we have two numbers multiplied together to get 0 The two numbers we multiply together are (x - 3) and (x + 2)

• By the Zero-Product Property either:

$$(x-3)=0$$

The first equation can be solved by adding 3 to each side to get:

$$(x+2)=0$$

We will start our discussion of Quadratic Equations with an example **Example:** Find the solution(s) to:

 $(x-3)\cdot(x+2)=0$

Here, we have two numbers multiplied together to get 0 The two numbers we multiply together are (x - 3) and (x + 2)

• By the Zero-Product Property either:

$$(x-3)=0$$

The first equation can be solved by adding 3 to each side to get: x = 3

$$(x+2)=0$$

We will start our discussion of Quadratic Equations with an example **Example:** Find the solution(s) to:

 $(x-3)\cdot(x+2)=0$

Here, we have two numbers multiplied together to get 0 The two numbers we multiply together are (x - 3) and (x + 2)

• By the Zero-Product Property either:

$$(x-3)=0$$

The first equation can be solved by adding 3 to each side to get: x = 3

$$(x+2)=0$$

The first equation can be solved by subtracting 2 from each side to get:

We will start our discussion of Quadratic Equations with an example **Example:** Find the solution(s) to:

 $(x-3)\cdot(x+2)=0$

Here, we have two numbers multiplied together to get 0 The two numbers we multiply together are (x - 3) and (x + 2)

• By the Zero-Product Property either:

$$(x-3)=0$$

The first equation can be solved by adding 3 to each side to get: x = 3

$$(x+2)=0$$

The first equation can be solved by subtracting 2 from each side to get:

$$x = -2$$

We will start our discussion of Quadratic Equations with an example **Example:** Find the solution(s) to:

 $(x-3)\cdot(x+2)=0$

Here, we have two numbers multiplied together to get 0 The two numbers we multiply together are (x - 3) and (x + 2)

• By the Zero-Product Property either:

$$(x-3)=0$$

The first equation can be solved by adding 3 to each side to get: x = 3

$$(x+2)=0$$

The first equation can be solved by subtracting 2 from each side to get:

$$\mathbf{x} = -2$$

Conclusion: The solutions to $(x - 3) \cdot (x + 2) = 0$ are: x = 3, -2