

# Types of solutions to Quadratic Equations

# Types of solutions to Quadratic Equations

Recall: The **▶ Quadratic Formula** says  $r_1$  and  $r_2$  are **▶ the roots** of:

$$ax^2 + bx + c = 0$$

# Types of solutions to Quadratic Equations

Recall: The **▶ Quadratic Formula** says  $r_1$  and  $r_2$  are **▶ the roots** of:

$$ax^2 + bx + c = 0$$
$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

# Types of solutions to Quadratic Equations

Recall: The [▶ Quadratic Formula](#) says  $r_1$  and  $r_2$  are [▶ the roots](#) of:

$$ax^2 + bx + c = 0$$
$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

We saw in [▶ Example 8](#) that if  $b^2 - 4ac = 0$  then we have one real solution of Multiplicity 2.

# Types of solutions to Quadratic Equations

Recall: The **▶ Quadratic Formula** says  $r_1$  and  $r_2$  are **▶ the roots** of:

$$ax^2 + bx + c = 0$$
$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

We saw in **▶ Example 8** that if  $b^2 - 4ac = 0$  then we have one real solution of Multiplicity 2.

We saw in **▶ Example 9** that if  $b^2 - 4ac < 0$  then we have no real solutions.

# Types of solutions to Quadratic Equations

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$ax^2 + bx + c = 0$$
$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

We saw in **Example 8** that if  $b^2 - 4ac = 0$  then we have one real solution of Multiplicity 2.

We saw in **Example 9** that if  $b^2 - 4ac < 0$  then we have no real solutions. If  $b^2 - 4ac > 0$  then we can compute the square root.

# Types of solutions to Quadratic Equations

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$ax^2 + bx + c = 0$$
$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

We saw in **Example 8** that if  $b^2 - 4ac = 0$  then we have one real solution of Multiplicity 2.

We saw in **Example 9** that if  $b^2 - 4ac < 0$  then we have no real solutions. If  $b^2 - 4ac > 0$  then we can compute the square root.

Since  $\sqrt{b^2 - 4ac} > 0$  adding and subtracting this number will give two distinct real solutions.

# Types of solutions to Quadratic Equations

Recall: The **▶ Quadratic Formula** says  $r_1$  and  $r_2$  are **▶ the roots** of:

$$ax^2 + bx + c = 0$$
$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

We saw in **▶ Example 8** that if  $b^2 - 4ac = 0$  then we have one real solution of Multiplicity 2.

We saw in **▶ Example 9** that if  $b^2 - 4ac < 0$  then we have no real solutions. If  $b^2 - 4ac > 0$  then we can compute the square root.

Since  $\sqrt{b^2 - 4ac} > 0$  adding and subtracting this number will give two distinct real solutions.

This means that we can determine what type of solutions we get from a quadratic equation based on  $b^2 - 4ac$



# Types of solutions to Quadratic Equations

Recall: The **▶ Quadratic Formula** says  $r_1$  and  $r_2$  are **▶ the roots** of:

$$ax^2 + bx + c = 0$$
$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

We saw in **▶ Example 8** that if  $b^2 - 4ac = 0$  then we have one real solution of Multiplicity 2.

We saw in **▶ Example 9** that if  $b^2 - 4ac < 0$  then we have no real solutions. If  $b^2 - 4ac > 0$  then we can compute the square root.

Since  $\sqrt{b^2 - 4ac} > 0$  adding and subtracting this number will give two distinct real solutions.

This means that we can determine what type of solutions we get from a quadratic equation based on  $b^2 - 4ac$

Because  $b^2 - 4ac$  plays an important role in our solutions, it gets a special name.

# Types of solutions to Quadratic Equations

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$ax^2 + bx + c = 0$$
$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

We saw in **Example 8** that if  $b^2 - 4ac = 0$  then we have one real solution of Multiplicity 2.

We saw in **Example 9** that if  $b^2 - 4ac < 0$  then we have no real solutions. If  $b^2 - 4ac > 0$  then we can compute the square root.

Since  $\sqrt{b^2 - 4ac} > 0$  adding and subtracting this number will give two distinct real solutions.

This means that we can determine what type of solutions we get from a quadratic equation based on  $b^2 - 4ac$

Because  $b^2 - 4ac$  plays an important role in our solutions, it gets a special name.

We call  $b^2 - 4ac$  *the discriminant*.