

Solving the quadratic equation:  $2x^2 - 3 = 0$ :

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The [Quadratic Formula](#) says  $r_1$  and  $r_2$  are [the roots](#) of:

$$ax^2 + bx + c = 0$$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$r_{1,2} = \frac{ax^2 + bx + c = 0}{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$r_{1,2} = \frac{ax^2 + bx + c = 0}{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$ax^2 + bx + c = 0$$
$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$
$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}}{2 \cdot 2}$$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$\begin{aligned} r_{1,2} &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ r_{1,2} &= \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}}{2 \cdot 2} \\ &= \frac{\pm \sqrt{0 + 24}}{4} \end{aligned}$$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$\begin{aligned} & ax^2 + bx + c = 0 \\ r_{1,2} &= \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a} \\ r_{1,2} &= \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}}{2 \cdot 2} \\ &= \frac{\pm \sqrt{0 + 24}}{4} \\ &= \frac{\pm \sqrt{24}}{4} \end{aligned}$$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$\begin{aligned} & ax^2 + bx + c = 0 \\ r_{1,2} &= \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a} \\ r_{1,2} &= \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}}{2 \cdot 2} \\ &= \frac{\pm \sqrt{0 + 24}}{4} \\ &= \frac{\pm \sqrt{24}}{4} \end{aligned}$$

The solutions to  $2x^2 - 3 = 0$  are:  $x = r_{1,2} = \frac{\pm \sqrt{24}}{4}$



## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$\begin{aligned} r_{1,2} &= \frac{ax^2 + bx + c = 0}{-(b) \pm \sqrt{(b)^2 - 4ac}} \\ r_{1,2} &= \frac{2a}{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}} \\ &= \frac{\pm \sqrt{0 + 24}}{2 \cdot 2} \\ &= \frac{4}{\pm \sqrt{24}} \\ &= \frac{\pm \sqrt{24}}{4} \end{aligned}$$

The solutions to  $2x^2 - 3 = 0$  are:  $x = r_{1,2} = \frac{\pm \sqrt{24}}{4}$

We can make this nicer if we notice:  $\sqrt{12} = \sqrt{4 \cdot 6}$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$\begin{aligned} r_{1,2} &= \frac{ax^2 + bx + c = 0}{-(b) \pm \sqrt{(b)^2 - 4ac}} \\ r_{1,2} &= \frac{2a}{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}} \\ &= \frac{\pm \sqrt{0 + 24}}{2 \cdot 2} \\ &= \frac{4}{\pm \sqrt{24}} \\ &= \frac{\pm \sqrt{24}}{4} \end{aligned}$$

The solutions to  $2x^2 - 3 = 0$  are:  $x = r_{1,2} = \frac{\pm \sqrt{24}}{4}$

We can make this nicer if we notice:  $\sqrt{12} = \sqrt{4 \cdot 6} = \sqrt{4} \cdot \sqrt{6}$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$\begin{aligned} r_{1,2} &= \frac{ax^2 + bx + c = 0}{-(b) \pm \sqrt{(b)^2 - 4ac}} \\ r_{1,2} &= \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}}{2 \cdot 2} \\ &= \frac{\pm \sqrt{0 + 24}}{4} \\ &= \frac{\pm \sqrt{24}}{4} \end{aligned}$$

The solutions to  $2x^2 - 3 = 0$  are:  $x = r_{1,2} = \frac{\pm \sqrt{24}}{4}$

We can make this nicer if we notice:  $\sqrt{12} = \sqrt{4 \cdot 6} = \sqrt{4} \cdot \sqrt{6} = 2\sqrt{6}$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$\begin{aligned} r_{1,2} &= \frac{ax^2 + bx + c = 0}{-(b) \pm \sqrt{(b)^2 - 4ac}} \\ r_{1,2} &= \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}}{2 \cdot 2} \\ &= \frac{\pm \sqrt{0 + 24}}{4} \\ &= \frac{\pm \sqrt{24}}{4} \end{aligned}$$

The solutions to  $2x^2 - 3 = 0$  are:  $x = r_{1,2} = \frac{\pm \sqrt{24}}{4}$

We can make this nicer if we notice:  $\sqrt{12} = \sqrt{4 \cdot 6} = \sqrt{4} \cdot \sqrt{6} = 2\sqrt{6}$

$$r_{1,2} = \frac{\pm 2\sqrt{6}}{4}$$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$\begin{aligned} r_{1,2} &= \frac{ax^2 + bx + c = 0}{-(b) \pm \sqrt{(b)^2 - 4ac}} \\ r_{1,2} &= \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}}{2 \cdot 2} \\ &= \frac{\pm \sqrt{0 + 24}}{4} \\ &= \frac{\pm \sqrt{24}}{4} \end{aligned}$$

The solutions to  $2x^2 - 3 = 0$  are:  $x = r_{1,2} = \frac{\pm \sqrt{24}}{4}$

We can make this nicer if we notice:  $\sqrt{12} = \sqrt{4 \cdot 6} = \sqrt{4} \cdot \sqrt{6} = 2\sqrt{6}$

$$\begin{aligned} r_{1,2} &= \frac{\pm 2\sqrt{6}}{4} \\ &= \frac{\pm \sqrt{6}}{2} \end{aligned}$$

## Solving the quadratic equation: $2x^2 - 3 = 0$ :

Recall: The **Quadratic Formula** says  $r_1$  and  $r_2$  are **the roots** of:

$$\begin{aligned} r_{1,2} &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ r_{1,2} &= \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 2 \cdot -3}}{2 \cdot 2} \\ &= \frac{\pm \sqrt{0 + 24}}{4} \\ &= \frac{\pm \sqrt{24}}{4} \end{aligned}$$

The solutions to  $2x^2 - 3 = 0$  are:  $x = r_{1,2} = \frac{\pm \sqrt{24}}{4}$

We can make this nicer if we notice:  $\sqrt{12} = \sqrt{4 \cdot 6} = \sqrt{4} \cdot \sqrt{6} = 2\sqrt{6}$

$$\begin{aligned} r_{1,2} &= \frac{\pm 2\sqrt{6}}{4} \\ &= \frac{\pm \sqrt{6}}{2} \end{aligned}$$

The solutions to  $2x^2 - 3 = 0$  are:  $x = r_{1,2} = \frac{\pm \sqrt{6}}{2}$