Solving the quadratic equation: $x^2 + 1 = 0$:

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ▸ Quadratic Formula says r_1 and r_2 are ▸ the roots of:

$$ax^2 + bx + c = 0$$

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ▸ Quadratic Formula says r_1 and r_2 are ▸ the roots of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ▸ Quadratic Formula says r_1 and r_2 are ▸ the roots of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ▸ Quadratic Formula says r_1 and r_2 are ▸ the roots of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ► Quadratic Formula says r_1 and r_2 are ► the roots of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$= \frac{\pm \sqrt{0 - 4}}{2}$$

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ▸ Quadratic Formula says r_1 and r_2 are ▸ the roots of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$= \frac{\pm \sqrt{0 - 4}}{2}$$

$$= \frac{\pm \sqrt{-4}}{2}$$

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ▸ Quadratic Formula says r_1 and r_2 are ▸ the roots of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$= \frac{\pm \sqrt{0 - 4}}{2}$$

$$= \frac{\pm \sqrt{-4}}{2}$$

$\sqrt{-4}$ is not a real number, so there are no real solutions!

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ⊳ Quadratic Formula says r_1 and r_2 are ⊳ the roots of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$= \frac{\pm \sqrt{0 - 4}}{2}$$

$$= \frac{\pm \sqrt{-4}}{2}$$

$\sqrt{-4}$ is not a real number, so there are no real solutions!

Conclusion: There are no real solutions to: $x^2 + 1 = 0$

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ▸ Quadratic Formula says r_1 and r_2 are ▸ the roots of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$= \frac{\pm \sqrt{0 - 4}}{2}$$

$$= \frac{\pm \sqrt{-4}}{2}$$

$\sqrt{-4}$ is not a real number, so there are no real solutions!

Conclusion: There are no real solutions to: $x^2 + 1 = 0$

Note 1: The reason there are no real solutions is that $-4 < 0$, so the square root is not a real number

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ▸ Quadratic Formula says r_1 and r_2 are ▸ the roots of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$= \frac{\pm \sqrt{0 - 4}}{2}$$

$$= \frac{\pm \sqrt{-4}}{2}$$

$\sqrt{-4}$ is not a real number, so there are no real solutions!

Conclusion: There are no real solutions to: $x^2 + 1 = 0$

Note 1: The reason there are no real solutions is that $-4 < 0$, so the square root is not a real number

Having $-4 < 0$ is because $-4 = b^2 - 4ac < 0$

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The (▸ Quadratic Formula) says r_1 and r_2 are (▸ the roots) of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$= \frac{\pm \sqrt{0 - 4}}{2}$$

$$= \frac{\pm \sqrt{-4}}{2}$$

$\sqrt{-4}$ is not a real number, so there are no real solutions!

Conclusion: There are no real solutions to: $x^2 + 1 = 0$

Note 1: The reason there are no real solutions is that $-4 < 0$, so the square root is not a real number

Having $-4 < 0$ is because $-4 = b^2 - 4ac < 0$

This is similar to what we say for repeated solutions!

Solving the quadratic equation: $x^2 + 1 = 0$:

Recall: The ⟨▸ Quadratic Formula⟩ says r_1 and r_2 are ⟨▸ the roots⟩ of:

$$ax^2 + bx + c = 0$$

$$r_{1,2} = \frac{-(b) \pm \sqrt{(b)^2 - 4ac}}{2a}$$

$$r_{1,2} = \frac{-(0) \pm \sqrt{(0)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1}$$

$$= \frac{\pm \sqrt{0 - 4}}{2}$$

$$= \frac{\pm \sqrt{-4}}{2}$$

$\sqrt{-4}$ is not a real number, so there are no real solutions!

Conclusion: There are no real solutions to: $x^2 + 1 = 0$

Note 1: The reason there are no real solutions is that $-4 < 0$, so the square root is not a real number

Having $-4 < 0$ is because $-4 = b^2 - 4ac < 0$

This is similar to what we say for repeated solutions!

Note 2: We will learn more about square roots of negatives soon!