

## Revisiting Factoring Quadratics - Example 3

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

We found that the general factored form will be:

$$6x^2 + x - 2 = a(x - r_1) \cdot (x - r_2)$$

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

We found that the general factored form will be:

$$6x^2 + x - 2 = a(x - r_1) \cdot (x - r_2)$$

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

We found that the general factored form will be:

$$6x^2 + x - 2 = a(x - r_1) \cdot (x - r_2)$$

We know that  $a = 6$  and we can find  $r_1, r_2$  using [The Quadratic Formula](#)

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

We found that the general factored form will be:

$$6x^2 + x - 2 = a(x - r_1) \cdot (x - r_2)$$

We know that  $a = 6$  and we can find  $r_1, r_2$  using [The Quadratic Formula](#)

► We can find the roots to be:

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

We found that the general factored form will be:

$$6x^2 + x - 2 = a(x - r_1) \cdot (x - r_2)$$

We know that  $a = 6$  and we can find  $r_1, r_2$  using [The Quadratic Formula](#)

[We can find](#) the roots to be:  $r_1, r_2 = \frac{1}{2}, \frac{-2}{3}$

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

We found that the general factored form will be:

$$6x^2 + x - 2 = a(x - r_1) \cdot (x - r_2)$$

We know that  $a = 6$  and we can find  $r_1, r_2$  using [The Quadratic Formula](#)

► We can find the roots to be:  $r_1, r_2 = \frac{1}{2}, -\frac{2}{3}$

So, we get the factored form:

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

We found that the general factored form will be:

$$6x^2 + x - 2 = a(x - r_1) \cdot (x - r_2)$$

We know that  $a = 6$  and we can find  $r_1, r_2$  using [The Quadratic Formula](#)

► We can find the roots to be:  $r_1, r_2 = \frac{1}{2}, \frac{-2}{3}$

So, we get the factored form:

$$6x^2 + x - 2 = 6\left(x - \frac{1}{2}\right) \cdot \left(x - \frac{-2}{3}\right)$$

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

We found that the general factored form will be:

$$6x^2 + x - 2 = a(x - r_1) \cdot (x - r_2)$$

We know that  $a = 6$  and we can find  $r_1, r_2$  using [The Quadratic Formula](#)

► We can find the roots to be:  $r_1, r_2 = \frac{1}{2}, \frac{-2}{3}$

So, we get the factored form:

$$6x^2 + x - 2 = 6\left(x - \frac{1}{2}\right) \cdot \left(x - \frac{-2}{3}\right)$$

If we wish, we can simplify:  $-\frac{-2}{3} = \frac{2}{3}$

## Revisiting Factoring Quadratics - Example 3

**Example:** Write  $6x^2 + x - 2$  in factored form:

We found that the general factored form will be:

$$6x^2 + x - 2 = a(x - r_1) \cdot (x - r_2)$$

We know that  $a = 6$  and we can find  $r_1, r_2$  using [The Quadratic Formula](#)

► We can find the roots to be:  $r_1, r_2 = \frac{1}{2}, \frac{-2}{3}$

So, we get the factored form:

$$6x^2 + x - 2 = 6\left(x - \frac{1}{2}\right) \cdot \left(x - \frac{-2}{3}\right)$$

If we wish, we can simplify:  $-\frac{-2}{3} = \frac{2}{3}$

$$6x^2 + x - 2 = 6\left(x - \frac{1}{2}\right) \cdot \left(x + \frac{2}{3}\right)$$