
Suppose we arrive at a gas station with 2 gallons of gas in the tank, and we fill it to 14 gallons.

Suppose we arrive at a gas station with 2 gallons of gas in the tank, and we fill it to 14 gallons. We can find how much the amount of gas changed by taking


the difference: 14 - 2 = 12

Suppose we arrive at a gas station with 2 gallons of gas in the tank, and we fill it to 14 gallons. We can find how much the amount of gas changed by taking the difference: 14 - 2 = 12In general, *Change* = end value - starting value


Suppose we arrive at a gas station with 2 gallons of gas in the tank, and we fill it to 14 gallons. We can find how much the amount of gas changed by taking the difference: 14 - 2 = 12In general, *Change* = *end value* - *starting value* What about change between points from (x_1, y_1) to (x_2, y_2) ?



Suppose we arrive at a gas station with 2 gallons of gas in the tank, and we fill it to 14 gallons. We can find how much the amount of gas changed by taking the difference: 14 - 2 = 12In general, *Change* = end value – starting value What about change between points from (x_1, y_1) to (x_2, y_2) ? One way we can look at the change between points is by looking at how much x and y each change individually.




Suppose we arrive at a gas station with 2 gallons of gas in the tank, and we fill it to 14 gallons. We can find how much the amount of gas changed by taking the difference: 14 - 2 = 12In general, *Change* = end value – starting value What about change between points from (x_1, y_1) to (x_2, y_2) ? One way we can look at the change between points is by looking at how much x and y each change individually. *Change in* x = end x value – starting x value




Suppose we arrive at a gas station with 2 gallons of gas in the tank, and we fill it to 14 gallons. We can find how much the amount of gas changed by taking the difference: 14 - 2 = 12In general, *Change* = end value – starting value What about change between points from (x_1, y_1) to (x_2, y_2) ? One way we can look at the change between points is by looking at how much x and y each change individually. *Change in* x = end x value – starting x value

Change in y = end y value – starting y value



Suppose we arrive at a gas station with 2 gallons of gas in the tank, and we fill it to 14 gallons. We can find how much the amount of gas changed by taking the difference: 14 - 2 = 12In general, *Change* = end value – starting value What about change between points from (x_1, y_1) to (x_2, y_2) ? One way we can look at the change between points is by looking at how much x and y each change individually. *Change in* x = end x value – starting x value $\Delta x = x_2 - x_1$ *Change in* y = end y value – starting y value



Suppose we arrive at a gas station with 2 gallons of gas in the tank, and we fill it to 14 gallons. We can find how much the amount of gas changed by taking the difference: 14 - 2 = 12In general, *Change* = end value – starting value What about change between points from (x_1, y_1) to (x_2, y_2) ? One way we can look at the change between points is by looking at how much x and y each change individually. *Change in* x = end x value – starting x value $\Delta x = x_2 - x_1$ *Change in* y = end y value – starting y value $\Delta y = y_2 - y_1$

