
Solving the quadratic equation: $x^2 + 3x + 3 = 0$:

Solving the quadratic equation: $x^2 + 3x + 3 = 0$:

Using the • Quadratic Formula

we get:



Solving the quadratic equation: $x^2 + 3x + 3 = 0$: Using the Quadratic Formula we get: $r_{1,2} = \frac{-(3) \pm \sqrt{(3)^2 - 4 \cdot 1 \cdot 3}}{\frac{2 \cdot 1}{2}}$ $= \frac{-3 \pm \sqrt{9 - 12}}{2}$

Solving the quadratic equation: $x^2 + 3x + 3 = 0$: Using the Quadratic Formula we get: $r_{1,2} = \frac{-(3) \pm \sqrt{(3)^2 - 4 \cdot 1 \cdot 3}}{\frac{2 \cdot 1}{2}}$ $= \frac{-3 \pm \sqrt{9 - 12}}{\frac{2}{2}}$ $= \frac{-3 \pm \sqrt{-3}}{2}$

Solving the quadratic equation: $x^2 + 3x + 3 = 0$: Using the • Quadratic Formula we get: $r_{1,2} = \frac{-(3) \pm \sqrt{(3)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1}$ $=\frac{-3\pm\sqrt{9-12}}{2}$ $= \frac{-3 \pm \sqrt[7]{-3}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot \sqrt{-1}}{2}$

Solving the quadratic equation: $x^2 + 3x + 3 = 0$: Using the • Quadratic Formula we get: $r_{1,2} = \frac{-(3) \pm \sqrt{(3)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1}$ $=\frac{-3\pm\sqrt{9-12}}{2}$ $=\frac{-3\pm\sqrt{-3}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot \sqrt{-1}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot i}{2}$

Solving the quadratic equation: $x^2 + 3x + 3 = 0$: Using the • Quadratic Formula we get: $r_{1,2} = \frac{-(3) \pm \sqrt{(3)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1}$ $=\frac{-3\pm\sqrt{9-12}}{2}$ $=\frac{-3\pm\sqrt{-3}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot \sqrt{-1}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot i}{2}$ **Conclusion:** The solutions to $x^2 + 3x + 3 = 0$ are the complex: $\frac{-3}{2} + \frac{\sqrt{3}}{2} \cdot i$ and $\frac{-3}{2} - \frac{\sqrt{3}}{2} \cdot i$

Solving the quadratic equation: $x^2 + 3x + 3 = 0$: Using the • Quadratic Formula we get: $r_{1,2} = \frac{-(3) \pm \sqrt{(3)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1}$ $=\frac{-3\pm\sqrt{9-12}}{2}$ $=\frac{-3\pm\sqrt{-3}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot \sqrt{-1}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot i}{2}$ **Conclusion:** The solutions to $x^2 + 3x + 3 = 0$ are the complex: $\frac{-3}{2} + \frac{\sqrt{3}}{2} \cdot i$ and $\frac{-3}{2} - \frac{\sqrt{3}}{2} \cdot i$ Note: The two solutions are complex conjugates

Solving the quadratic equation: $x^2 + 3x + 3 = 0$: Using the • Quadratic Formula we get: $r_{1,2} = \frac{-(3) \pm \sqrt{(3)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1}$ $=\frac{-3\pm\sqrt{9-12}}{2}$ $=\frac{-3\pm\sqrt{-3}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot \sqrt{-1}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot i}{2}$ **Conclusion:** The solutions to $x^2 + 3x + 3 = 0$ are the complex: $\frac{-3}{2} + \frac{\sqrt{3}}{2} \cdot i$ and $\frac{-3}{2} - \frac{\sqrt{3}}{2} \cdot i$ Note: The two solutions are complex conjugates This is not a coincidence. The difference between the two solutions is the \pm before the imaginary part with i

Solving the quadratic equation: $x^2 + 3x + 3 = 0$: Using the • Quadratic Formula we get: $r_{1,2} = \frac{-(3) \pm \sqrt{(3)^2 - 4 \cdot 1 \cdot 3}}{2 \cdot 1}$ $=\frac{-3\pm\sqrt{9-12}}{2}$ $=\frac{-3\pm\sqrt{-3}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot \sqrt{-1}}{2}$ $= \frac{-3 \pm \sqrt{3} \cdot i}{2}$ **Conclusion:** The solutions to $x^2 + 3x + 3 = 0$ are the complex: $\frac{-3}{2} + \frac{\sqrt{3}}{2} \cdot i$ and $\frac{-3}{2} - \frac{\sqrt{3}}{2} \cdot i$ Note: The two solutions are complex conjugates This is not a coincidence. The difference between the two solutions is the \pm before the imaginary part with i **Conclusion:** In general, if a + bi is a solution to a quadratic equation, then its complex conjugate a - bi is a solution, also.