

## Powers of $i$ - Examples

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:  
 $i^m = i^n$  if  $m - n$  is a multiple of 4.

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16}$

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4$  because  $16 - 4 = 12$  is a multiple of 4

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4



## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18}$

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2$  because  $18 - 2 = 16$  is a multiple of 4

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

**Example:**  $i^{47}$

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

**Example:**  $i^{47} = i^7$  because  $47 - 7 = 40$  is a multiple of 4

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

**Example:**  $i^{47} = i^7 = i^3$  because  $7 - 3 = 4$  is a multiple of 4



## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

**Example:**  $i^{47} = i^7 = i^3 = -i$  because  $7 - 3 = 4$  is a multiple of 4

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

**Example:**  $i^{47} = i^7 = i^3 = -i$  because  $7 - 3 = 4$  is a multiple of 4

If we want to do this all in one step,  $i^m = i^r$  if  $r$  is the remainder when we divide  $m$  by 4

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

**Example:**  $i^{47} = i^7 = i^3 = -i$  because  $7 - 3 = 4$  is a multiple of 4

If we want to do this all in one step,  $i^m = i^r$  if  $r$  is the remainder when we divide  $m$  by 4

**Example:**  $i^{522}$

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

**Example:**  $i^{47} = i^7 = i^3 = -i$  because  $7 - 3 = 4$  is a multiple of 4

If we want to do this all in one step,  $i^m = i^r$  if  $r$  is the remainder when we divide  $m$  by 4

**Example:**  $i^{522}$

Dividing 522 by 4 we get a remainder  $r = 2$

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

**Example:**  $i^{47} = i^7 = i^3 = -i$  because  $7 - 3 = 4$  is a multiple of 4

If we want to do this all in one step,  $i^m = i^r$  if  $r$  is the remainder when we divide  $m$  by 4

**Example:**  $i^{522} = i^2$

Dividing 522 by 4 we get a remainder  $r = 2$

## Powers of $i$ - Examples

► We saw that powers of  $i$  follow the pattern:

$i^m = i^n$  if  $m - n$  is a multiple of 4.

And also computed:

$$i^1 = i$$

$$i^2 = -1$$

$$i^3 = -i$$

$$i^4 = 1$$

Using these, we can compute  $i$  to any integer power.

**Example:**  $i^{16} = i^4 = 1$  because  $16 - 4 = 12$  is a multiple of 4

**Example:**  $i^{18} = i^2 = -1$  because  $18 - 2 = 16$  is a multiple of 4

For larger powers, it gets harder to figure out how to reduce

It can be helpful to reduce in multiple steps

**Example:**  $i^{47} = i^7 = i^3 = -i$  because  $7 - 3 = 4$  is a multiple of 4

If we want to do this all in one step,  $i^m = i^r$  if  $r$  is the remainder when we divide  $m$  by 4

**Example:**  $i^{522} = i^2 = -1$

Dividing 522 by 4 we get a remainder  $r = 2$